

Policies Repository



Policy Title Adalimumab (Humira®)

Policy Number FS.CLIN.22

Application of Pharmacy Policy is determined by benefits and contracts. Benefits may vary based on product line, group or contract. Some medications may be subject to precertification, age, gender or quantity edits. Individual member benefits must be verified.

This Pharmacy Policy document describes the status of pharmaceutical information and/or technology at the time the document was developed. Since that time, new information relating to drug efficacy, interactions, contraindications, dosage, administration routes, safety or FDA approval may have changed. If the Medical/Pharmacy Reviewer is aware of any new information on the subject of this document, please provide it promptly to the Medical/Pharmacy Policy Department. This information may include new FDA approved indications, withdrawals or other FDA alerts. This type of information is relevant not only when considering whether this Policy should be updated, but also when applying it to current requests for coverage.

Members are advised to use participating pharmacies in order to receive the highest level of benefits.

Policy **Adalimumab (Humira®)** is indicated for the treatment of crohn's disease, moderate to severe rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, moderate to severe juvenile idiopathic arthritis (JIA) and moderate to severe plaque psoriasis.

The use of adalimumab (Humira®) requires prior authorization (ie, clinical pharmacy and/or Medical Director review).

Policy Description **Adalimumab (Humira®)** is a recombinant human IgG1 monoclonal antibody that blocks tumor necrosis factor (TNF). TNF is a naturally occurring cytokine that plays a role in inflammatory and immune responses. Elevated TNF levels can cause inflammation and joint destruction in individuals with rheumatoid arthritis, ankylosing spondylitis, juvenile idiopathic arthritis, and psoriatic arthritis. Adalimumab (Humira®) binds to TNF-alpha, blocks its interaction with surface TNF receptors, and causes surface lysis (breakdown), leading to decreased inflammation and joint destruction. Adalimumab (Humira®) does not bind to or inactivate TNF-beta. Adalimumab (Humira®) also modulates biological responses that are induced or regulated by TNF, including changes in the levels of adhesion molecules responsible for leukocyte migration. Adalimumab (Humira®) decreases C-reactive protein, erythrocyte sedimentation rate, IL-6, and matrix metalloproteinases MMP-1 and MMP-3.

Policy Guideline Inclusion **Adalimumab (Humira®)** is approved when **any** of the following inclusion criteria is met:

- Documentation of a diagnosis of moderate to severe Rheumatoid Arthritis, Ankylosing Spondylitis or Psoriatic Arthritis and ALL of the following:
 - Patient is an adult (\geq 18 years)
 - Medication is being recommended and prescribed by a rheumatologist
 - Patient had at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) OR contraindication to ALL of the following DMARDs:

- Methotrexate
 - Hydroxychloroquine
 - Leflunomide
 - Azathioprine
 - Sulfasalazine
 - Etanercept (Enbrel®)
 - Patient is not on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
 - Patient does not have active infections or sepsis
 - Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
 - Patient does not have active malignancy
- Documentation of a diagnosis of moderate to severe Juvenile Idiopathic Arthritis (JIA) and ALL of the following:
 - Patient is ≥ 4 years old
 - Medication is being recommended and prescribed by a rheumatologist
 - Patient had at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) OR contraindication to ALL of the following DMARDs:
 - Methotrexate
 - Hydroxychloroquine
 - Leflunomide
 - Azathioprine
 - Sulfasalazine
 - Etanercept (Enbrel®)
 - Patient is not on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
 - Patient does not have active infections or sepsis
 - Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
 - Patient does not have active malignancy
- Documentation of a diagnosis of moderate to severe chronic Plaque Psoriasis and ALL of the following:
 - Patient is an adult (≥ 18 years)
 - Medication is being recommended and prescribed by a dermatologist
 - Patient had at least a 30 day trial and failure with ONE of the following drugs OR contraindication to ALL of the following drugs:
 - Topical Calcipotriene containing products
 - Topical Anthralin
 - Topical Steroids
 - Topical immunomodulators (Elidel®, Protopic®)
 - Topical retinoids
 - Efudex
 - Etanercept (Enbrel®)
 - Patient is not on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
 - Patient does not have active infections or sepsis
 - Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
 - Patient does not have active malignancy

- Documentation of a diagnosis of Crohn's Disease and ALL of the following:
 - Patient is an adult (≥ 18 years old)
 - Medication is being recommended and prescribed by a gastroenterologist
 - Patient had at least a 30 day trial and failure with Infliximab (Remicade[®]) OR at least a 30 day trial and failure with one drug from any TWO of the following groups OR contraindication to ALL of the following groups:
 - Corticosteroids: Budesonide (Entocort[®] EC), Prednisone, Hydrocortisone, Methylprednisolone
 - Aminosalicylates: Sulfasalazine, Mesalamine (Asacol[®], Rowasa[®], Canasa[®], Pentasa[®]), Olsalazine (Dipentum[®]), Balsalazide (Colazal[™])
 - Immunomodulators: Azathioprine, 6-mercaptopurine, Cyclosporine, Sirolimus (Prograf[®]), Methotrexate
 - Antibiotics: Metronidazole or Fluoroquinolones
 - Patient is not on concurrent therapy with Anakinra (Kineret[®]) or other tumor necrosis factor antagonists
 - Patient does not have active infections or sepsis
 - Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
 - Patient does not have active malignancy

Policy Guideline Exclusion

For a diagnosis of moderate to severe Rheumatoid Arthritis, Ankylosing Spondylitis or Psoriatic Arthritis, **Adalimumab (Humira[®])** is denied when **any** of the following exclusion criteria is present:

- Patient is less than 18 years old
- Medication is not being prescribed by a rheumatologist
- Patient does not have at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) or contraindication to ALL of the following DMARDs:
 - Methotrexate
 - Hydroxychloroquine
 - Leflunomide
 - Azathioprine
 - Sulfasalazine
 - Etanercept (Enbrel[®])
- Patient is on concurrent therapy with Anakinra (Kineret[®]) or other tumor necrosis factor antagonists
- Patient has active infections or sepsis
- Patient has not been evaluated using tuberculin skin test
- Patient has active or latent tuberculosis
- Patient has active malignancy

For a diagnosis of moderate to severe Juvenile Idiopathic Arthritis (JIA), **Adalimumab (Humira[®])** is denied when **any** of the following exclusion criteria is present:

- Patient is less than 4 years old
- Medication is not being prescribed by a rheumatologist
- Patient does not have at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) or contraindication to ALL of the following DMARDs:
 - Methotrexate
 - Hydroxychloroquine

- Leflunomide
- Azathioprine
- Sulfasalazine
- Etanercept (Enbrel®)
- Patient is on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
- Patient has active infections or sepsis
- Patient has not been evaluated using tuberculin skin test
- Patient has active or latent tuberculosis
- Patient has active malignancy

For a diagnosis of moderate to severe chronic Plaque Psoriasis, **Adalimumab (Humira®)** is denied when **any** of the following exclusion criteria is present:

- Patient is less than 18 years old
- Medication is not being prescribed by a dermatologist
- Patient does not have at least a 30 day trial and failure with ONE of the following drugs OR contraindication to ALL of the following drugs:
 - Topical Calcipotriene containing products
 - Topical Anthralin
 - Topical Steroids
 - Topical immunomodulators (Elidel®, Protopic®)
 - Topical retinoids
 - Efudex
 - Etanercept (Enbrel®)
- Patient is on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
- Patient has active infections or sepsis
- Patient has not been evaluated using tuberculin skin test
- Patient has active or latent tuberculosis
- Patient has active malignancy

For a diagnosis of Crohn's Disease, **adalimumab (Humira®)** is denied when **any** of the following exclusion criteria is present:

- Patient is less than 18 years old
- Medication is not being prescribed by a gastroenterologist
- Patient does not have at least a 30 day trial and failure with Infliximab (Remicade®) OR at least a 30 day trial and failure with one drug from any TWO of the following groups OR contraindication to ALL of the following groups:
 - Corticosteroids: Budesonide (Entocort® EC), Prednisone, Hydrocortisone, Methylprednisolone
 - Aminosalicylates: Sulfasalazine, Mesalamine (Asacol®, Rowasa®, Canasa®, Pentasa®), Olsalazine (Dipentum®), Balsalazide (Colazal™)
 - Immunomodulators: Azathioprine, 6-mercaptopurine, Cyclosporine, Sirolimus (Prograf®), Methotrexate
 - Antibiotics: Metronidazole or Fluoroquinolones
- Patient is on concurrent therapy with Anakinra (Kineret®) or other tumor necrosis factor antagonists
- Patient has active infections or sepsis
- Patient has not been evaluated using tuberculin skin test
- Patient has active or latent tuberculosis

- Patient has active malignancy

Policy List of Applicable Drugs

Brand Name	Generic Name
Humira	adalimumab

Dosing and Administration

Refer to the specific manufacturer's prescribing information for administration and dosage details, contraindications, and Black Box warnings.

Policy References

American College of Rheumatology Subcommittee on Rheumatoid Arthritis Guidelines. Guidelines for the management of rheumatoid arthritis: 2002 update. [American College of Rheumatology Web site]. Available at: <http://www.rheumatology.org/publications/guidelines/raguidelines02.asp?aud=mem>. Accessed July 1, 2009.

American Thoracic Society, Centers for Disease Control and Prevention. Targeted tuberculin testing and treatment of latent tuberculosis infection. *Am J Respir Crit Care Med*. 2000;161:S221-S247.

Barrera P, Joosten LA, den Broeder AA, et al: Effects of treatment with a fully human anti-tumour necrosis factor alpha monoclonal antibody on the local and systemic homeostasis of interleukin 1 and TNFalpha in patients with rheumatoid arthritis. *Ann Rheum Dis* 2001; 60:660-669.

Breedveld FC, Weisman MH, Kavanaugh AF, Cohen SB, Pavelka K, van Vollenhoven R, et al. The PREMIER study: A multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;54(1):26-37.

Colombel JF, Sandborn WJ, Rutgeerts P, et al: Adalimumab for maintenance of clinical response and remission in patients with Crohn's disease: the CHARM trial. *Gastroenterology* 2007; 132(1):52-65.

Donahue KE, Gartlehner G, Jonas DE, Lux LJ, Thieda P, Jonas B, Hansen RA, Morgan LC, Williams SC, Lohr KN. Comparative Effectiveness of Drug Therapy for Rheumatoid Arthritis and Psoriatic Arthritis in Adults. Comparative Effectiveness Review No. 11. (Prepared by RTI-University of North Carolina Evidence-based Practice Center under Contract No. 290-02-0016.) Rockville, MD: Agency for Healthcare Research and Quality. November 2007. Available at: <http://derp.ohsu.edu/about/final-products.cfm>.

Firestein GS, Boyle DL, Yu C, et al. Synovial interleukin-1 receptor antagonist and interleukin-1 balance in rheumatoid arthritis. *Arthritis Rheum*. 1994; 37:644-652.

Finckh A, Simard JF, Duryea J, Liang MH, Huang J, Daneel S, et al. The effectiveness of anti-tumor necrosis factor therapy in preventing progressive radiographic joint damage in rheumatoid arthritis: a population-based study. *Arthritis Rheum* 2006;54(1):54-9.

Fries JF, Spitz P, Kraines RG, Holman HR. Measurement of patient outcome in arthritis. *Arthritis Rheum*. 1980;23(2):137-145.

Furst DE, Schiff MH, Fleischmann RM, Strand V, Birbara CA, Compagnone D, et al. Adalimumab, a fully human anti tumor necrosis factor-alpha monoclonal antibody, and concomitant standard antirheumatic therapy for the treatment of rheumatoid arthritis: results of STAR (Safety Trial of Adalimumab in Rheumatoid Arthritis). *J Rheumatol* 2003;30(12):2563-71.

Gartlehner G, Hansen RA, Thieda P, Jonas B, Lohr KN, Carey T. Drug Class Review on Targeted Immunomodulators [Final Report] January 2007. Oregon Health & Science University. Drug

Effectiveness Review Project [available at <http://derp.ohsu.edu/about/final-products.cfm>]

Genant HK, Jiang Y, Peterfy C, Lu Y, Re ´ dei J, Countryman PJ. Assessment of rheumatoid arthritis using a modified scoring method on digitized and original radiographs. *Arthritis Rheum.* 1998;41(9):1583-1590.

Gladman DD. Traditional and newer therapeutic options for psoriatic arthritis: an evidence-based review. *Drugs* 2005;65(9):1223-38.

Gladman DD, Mease PJ, Cifaldi MA, et al: Adalimumab improves joint-related and skin-related functional impairment in patients with psoriatic arthritis: patient-reported outcomes of the Adalimumab Effectiveness in Psoriatic Arthritis Trial. *Ann Rheum Dis* 2007; 66(2):163-168.

Gordon KB, Langley RG, Leonardi C, et al: Clinical response to adalimumab treatment in patients with moderate to severe psoriasis: double-blind, randomized controlled trial and open-label extension study. *J Am Acad Dermatol* 2006; 55(4):598-606.

Humira[®] (Adalimumab) In: Drugdex [online through Micromedex Healthcare Series]. Greenwood Village, CO: Thomson Micromedex. Accessed July 1, 2009.

Humira[®] (Adalimumab) In: Facts and Comparisons [online through Facts and Comparisons Online]. Indy, IN: Walter Kluwer Health Inc. Accessed July 1, 2009.

Humira[®] (Adalimumab) [package insert]. North Chicago, IL: Abbott; 2007.

Kavanaugh AF, Ritchlin CT. Systematic review of treatments for psoriatic arthritis: an evidence based approach and basis for treatment guidelines. *J Rheumatol* 2006;33(7):1417-21.

Keystone E, Weinblatt M, Furst D, et al: The ARMADA trial: a double-blind placebo controlled trial of the fully human anti-TNF monoclonal antibody: adalimumab (D2E7), in patients with active RA on methotrexate (MTX) (abstract 965). *Arthritis Rheum* 2001a; 44(9):S213.

Mease PJ, Gladman DD, Ritchlin CT, et al: Adalimumab for the treatment of patients with moderately to severely active psoriatic arthritis: results of a double-blind, randomized, placebo-controlled trial. *Arthritis Rheum* 2005; 52(10):3279-3289.

National Institute for Clinical Excellence (NICE). A systematic review of the effectiveness of adalimumab, etanercept and infliximab for the treatment of rheumatoid arthritis in adults and an economic evaluation of their cost-effectiveness. London (UK): National Institute for Clinical Excellence (NICE); 2005 October.

Papadakis KA, Shaye OA, Vasiliauskas EA, et al: Safety and efficacy of adalimumab (D2E7) in Crohn's disease patients with an attenuated response to infliximab. *Am J Dig Dis* 2005; 100(1):75-79.

Pincus T, Summey JA, Soraci SA Jr, Wallston KA, Hummon NP. Assessment of patient satisfaction in activities of daily living using a modified Stanford Health Assessment Questionnaire. *Arthritis Rheum.* 1983;26(11):1346-1353.

Rau R, Herborn G, Sander O, et al: Long-term treatment with the fully human anti-TNF-antibody D2E7 slows radiographic disease progression in rheumatoid arthritis (abstract 1978). *Arthritis Rheum* 1999; 42(9 Suppl S):S400.

Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg L, Mariotto A, Feuer EF, Edwards BK (eds). *SEER Cancer Statistics Review, 1975-2001*, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2001/.

Scallon BJ, Moore MA, Trinh H, et al. Chimeric anti-TNF α monoclonal antibody cA2 binds recombinant transmembrane TNF α and activates immune effector functions. *Cytokine*. 1995;7:251-259.

Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active ulcerative colitis. A randomized study. *N Engl J Med*. 1987;317(26):1625-1629.

Targan SR, Hanauer SB, van Deventer SJH, et al. A short-term study of chimeric monoclonal antibody cA2 to tumor necrosis factor α for Crohn's disease. *N Engl J Med*. 1997;337(15):1029-1035.

Van der Heijde DM, van Leeuwen MA, van Riel PL, et al. Biannual radiographic assessments of hands and feet in a three-year prospective follow-up of patients with early rheumatoid arthritis. *Arthritis Rheum*. 1992;35(1):26-34.

Van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the New York criteria. *Arthritis Rheum*. 1984;27(4):361-368.

Ware JE Jr, Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) Project. *J Clin Epidemiol*. 1998;51(11):903-912.

Weinblatt ME, Keystone EC, Furst DE, Moreland LW, Weisman MH, Birbara CA, et al. Adalimumab, a fully human anti-tumor necrosis factor alpha monoclonal antibody, for the treatment of rheumatoid arthritis in patients taking concomitant methotrexate: the ARMADA trial. *Arthritis Rheum* 2003;48(1):35-45.

Weisman M, Keystone E, Paulus H, et al: A dose escalation study designed to demonstrate the safety, tolerability and efficacy of the fully human anti-TNF antibody, D2E7, given in combination with methotrexate (MTX) in patients with active RA (abstract 1948). *Arthritis Rheum* 2000; 43(9 Suppl S):S391.

Policy Link to Related Policies

Printed

09/01/2009 10:13:06

The Policy Bulletins on this web site were developed to assist AmeriHealth and its subsidiaries ("AmeriHealth") in administering the provisions of the respective benefit programs, and do not constitute a contract. If you are an AmeriHealth member, please refer to your specific benefit program for the terms, conditions, limitations and exclusions of your coverage. AmeriHealth does not provide health care services, medical advice or treatment, or guarantee the outcome or results of any medical services/treatments. The facility and professional providers are responsible for providing medical advice and treatment. Facility and professional providers are independent contractors and are not employees or agents of AmeriHealth. If you have a specific medical condition, please consult with your doctor. AmeriHealth reserves the right at any time to change or update its Policy Bulletins. ©2008 AmeriHealth, Inc. All Rights Reserved. Current Procedural Terminology ©2008 American Medical Association. All Rights Reserved.