

Policies Repository



Policy Title Anakinra (Kineret®)

Policy Number FS.CLIN.50

Application of Pharmacy Policy is determined by benefits and contracts. Benefits may vary based on product line, group or contract. Some medications may be subject to precertification, age, gender or quantity edits. Individual member benefits must be verified.

This Pharmacy Policy document describes the status of pharmaceutical information and/or technology at the time the document was developed. Since that time, new information relating to drug efficacy, interactions, contraindications, dosage, administration routes, safety or FDA approval may have changed. If the Medical/Pharmacy Reviewer is aware of any new information on the subject of this document, please provide it promptly to the Medical/Pharmacy Policy Department. This information may include new FDA approved indications, withdrawals or other FDA alerts. This type of information is relevant not only when considering whether this Policy should be updated, but also when applying it to current requests for coverage.

Members are advised to use participating pharmacies in order to receive the highest level of benefits.

Policy **Anakinra (Kineret®)** is indicated for the treatment of moderate to severe rheumatoid arthritis.

The use of anakinra (Kineret®) requires prior authorization (ie, clinical pharmacy and/or Medical Director review).

Policy Description

Anakinra (Kineret®) is a synthetic, injectable interleukin-1 (IL-1) receptor antagonist that blocks the effects of human IL-1. IL-1 is a protein produced by many cells in the body, but it is found in increased amounts within joints that are inflamed from rheumatoid arthritis (RA). IL-1 attaches to receptors on the tissues within and surrounding the joints, as well as to the cells that are responsible for inflammation (eg, white blood cells). The attachment of IL-1 to its receptors activates inflammation and the release of enzymes that destroy cartilage and bone, which contributes to joint pain and swelling. By binding to these receptors, anakinra (Kineret®) prevents the inflammatory and enzyme-releasing effects of IL-1, thereby reducing joint pain and swelling.

Policy Guideline Inclusion

Anakinra (Kineret®) is approved when **all** of the following inclusion criteria are met:

- Documented diagnosis of moderate to severe Rheumatoid Arthritis and ALL of the following:
 - Patient is an adult (≥ 18 years)
 - Medication is being recommended and prescribed by a rheumatologist
 - Patient had at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) OR contraindication to ALL of the following DMARDs:
 - Methotrexate
 - Hydroxychloroquine
 - Leflunomide
 - Azathioprine
 - Sulfasalazine

- Adalimumab (Humira®)
 - Etanercept (Enbrel®)
- Patient does not have concurrent therapy with tumor necrosis factor antagonists
- Patient does not have active infections or sepsis
- Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
- Patient does not have active malignancy

Policy Guideline Exclusion

Anakinra (Kineret®) is denied when the following exclusion criteria is met:

- No documented diagnosis of moderate to severe Rheumatoid Arthritis and ALL of the following:
 - Patient is an adult (≥ 18 years)
 - Medication is being recommended and prescribed by a rheumatologist
 - Patient had at least a 30 day trial and failure with ONE of the following disease-modifying anti-rheumatic drugs (DMARDs) OR contraindication to ALL of the following DMARDs:
 - Methotrexate
 - Hydroxychloroquine
 - Leflunomide
 - Azathioprine
 - Sulfasalazine
 - Adalimumab (Humira®)
 - Etanercept (Enbrel®)
 - Patient does not have concurrent therapy with tumor necrosis factor antagonists
 - Patient does not have active infections or sepsis
 - Patient has been evaluated (i.e. tuberculin skin test) and does not have active or latent tuberculosis
 - Patient does not have active malignancy

Policy List of Applicable Drugs

Brand Name	Generic Name
Kineret	anakinra

Dosing and Administration

Refer to the specific manufacturer's prescribing information for administration and dosage details, contraindications, and Black Box warnings.

Policy References

American College of Rheumatology Subcommittee on Rheumatoid Arthritis Guidelines. Guidelines for the management of rheumatoid arthritis: 2002 update. [American College of Rheumatology Web site]. Available at: <http://www.rheumatology.org/publications/guidelines/raguidelines02.asp?aud=mem>. Accessed February 28, 2008.

American Thoracic Society, Centers for Disease Control and Prevention. Targeted tuberculin testing and treatment of latent tuberculosis infection. Am J Respir Crit Care Med. 2000;161:S221-S247.

Firestein GS, Boyle DL, Yu C, et al. Synovial interleukin-1 receptor antagonist and interleukin-1 balance in rheumatoid arthritis. Arthritis Rheum. 1994; 37:644-652.

Fries JF, Spitz P, Kraines RG, Holman HR. Measurement of patient outcome in arthritis. Arthritis Rheum. 1980;23(2):137-145.

Gartlehner G, Hansen RA, Thieda P, Jonas B, Lohr KN, Carey T. Drug Class Review on Targeted Immunomodulators [Final Report] January 2007. Oregon Health & Science University. Drug

Effectiveness Review Project [available at
<http://www.ohsu.edu/drugeffectiveness/reports/final.cfm>]

Genant HK, Jiang Y, Peterfy C, Lu Y, Re ´dei J, Countryman PJ. Assessment of rheumatoid arthritis using a modified scoring method on digitized and original radiographs. *Arthritis Rheum.* 1998;41(9):1583-1590.

Kineret[®] (Anakinra) In: Facts and Comparisons [online through Facts and Comparisons Online]. Indy, IN: Walter Kluwer Health Inc. Accessed November 13, 2007.

Kineret[®] (Anakinra) In: Drugdex [online through Micromedex Healthcare Series]. Greenwood Village, CO: Thomson Micromedex. Accessed November 13, 2007.

Kineret[®] (Anakinra) [package insert]. Thousand Oaks, CA: Amgen[®]; 2006.

Pincus T, Summey JA, Soraci SA Jr, Wallston KA, Hummon NP. Assessment of patient satisfaction in activities of daily living using a modified Stanford Health Assessment Questionnaire. *Arthritis Rheum.* 1983;26(11):1346-1353.

Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg L, Mariotto A, Feuer EF, Edwards BK (eds). SEER Cancer Statistics Review, 1975-2001, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2001/.

Van der Heijde DM, van Leeuwen MA, van Riel PL, et al. Biannual radiographic assessments of hands and feet in a three-year prospective follow-up of patients with early rheumatoid arthritis. *Arthritis Rheum.* 1992;35(1):26-34.

Ware JE Jr, Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) Project. *J Clin Epidemiol.* 1998;51(11):903-912

Policy Link to Related Policies

Printed

11/11/2008 15:16:04

The Policy Bulletins on this web site were developed to assist AmeriHealth and its subsidiaries ("AmeriHealth") in administering the provisions of the respective benefit programs, and do not constitute a contract. If you are an AmeriHealth member, please refer to your specific benefit program for the terms, conditions, limitations and exclusions of your coverage. AmeriHealth does not provide health care services, medical advice or treatment, or guarantee the outcome or results of any medical services/treatments. The facility and professional providers are responsible for providing medical advice and treatment. Facility and professional providers are independent contractors and are not employees or agents of AmeriHealth. If you have a specific medical condition, please consult with your doctor. AmeriHealth reserves the right at any time to change or update its Policy Bulletins. ©2008 AmeriHealth, Inc. All Rights Reserved. Current Procedural Terminology ©2008 American Medical Association. All Rights Reserved.